


Amendments to the Claims:

This listing of claims will replace all prior versions and listing of claims in the application.

Listing of Claims:

Claim 1 (currently amended): A quinazoline derivative of the formula I:



I

wherein:

m is 0, 1 or 2;

each **R¹**, which may be the same or different, is selected from hydroxy, (1-6C)alkoxy,

(3-7C)cycloalkyl-oxy and (3-7C)cycloalkyl-(1-6C)alkoxy, and

wherein any CH₂ or CH₃ group within a **R¹** substituent optionally bears on each said CH₂ or CH₃ group one or more halogeno or (1-6C)alkyl substituents, or a substituent selected from

hydroxy and (1-6C)alkoxy;

R² is hydrogen or (1-4C)alkyl;

n is 0, 1, 2, 3 or 4;

each **R³**, which may be the same or different, is selected from cyano, halogeno, (1-4C)alkyl, trifluoromethyl, (1-4C)alkoxy, (2-4C)alkenyl and (2-4C)alkynyl;

X¹ is selected from O, S, SO, SO₂, N(R⁷), CH(OR⁷), CON(R⁷), N(R⁷)CO, SO₂N(R⁷), N(R⁷)SO₂, OC(R⁷)₂, C(R⁷)₂O, SC(R⁷)₂, C(R⁷)₂S, CO, C(R⁷)₂N(R⁷) and N(R⁷)C(R⁷)₂; wherein:

each **R⁷**, which may be the same or different, is hydrogen or (1-6C)alkyl;

Q¹ is aryl, or heteroaryl, and

wherein Q¹ optionally bears one or more substituents, which may be the same or different, selected from halogeno, cyano, nitro, hydroxy, amino, carboxy, carbamoyl, sulfamoyl, formyl, mercapto, (1-6C)(1-4C)alkyl, (2-8C)alkenyl, (2-8C)alkynyl, and (1-6C)(1-4C)alkoxy, (2-6C)alkenyloxy, (2-6C)alkynyloxy, (1-6C)alkylthio, (1-6C)alkylsulfinyl, (1-6C)alkylsulfonyl, (1-6C)alkylamino, di-[(1-6C)alkyl]amino, (1-6C)alkoxy-carbonyl, N-(1-6C)alkyl-carbamoyl, N,N-di-[(1-6C)alkyl]carbamoyl, (2-6C)alkanoyl, (3-6C)alkenoyl, (3-6C)alkynoyl, (2-6C)alkanoyloxy, (2-6C)alkanoylamino, N-(1-6C)alkyl-(2-6C)alkanoylamino, (3-6C)alkenoylamino, N-(1-6C)alkyl-(3-6C)alkenoylamino, (3-6C)alkynoylamino, N-(1-6C)alkyl-(3-6C)alkynoylamino, N-(1-6C)alkyl-sulfamoyl, N,N-di-[(1-6C)alkyl]sulfamoyl, (1-6C)alkanesulfonylamino, N-(1-6C)alkyl-(1-6C)alkanesulfonylamino, and a group of the formula: $-X^2-R^8$ wherein X² is a direct bond or is selected from O, CO and N(R⁹), wherein:

R⁹ is hydrogen or (1-6C)alkyl, and

R⁸ is halogeno (1-6C)alkyl, hydroxy (1-6C)alkyl, carboxy (1-6C)alkyl, (1-6C)alkoxy (1-6C)alkyl, cyano (1-6C)alkyl, amino (1-6C)alkyl, N-(1-6C)alkylamino (1-6C)alkyl, N,N-di-[(1-6C)alkyl]amino (1-6C)alkyl, (2-6C)alkanoylamino (1-6C)alkyl, N-(1-6C)alkyl-(2-6C)alkanoylamino (1-6C)alkyl, (1-6C)alkoxy-carbonylamino (1-6C)alkyl, carbamoyl (1-6C)alkyl, N-(1-6C)alkyl-carbamoyl (1-6C)alkyl, N,N-di-[(1-6C)alkyl]carbamoyl (1-6C)alkyl, (1-6C)alkylthio (1-6C)alkyl, (1-6C)alkylsulfinyl (1-6C)alkyl, (1-6C)alkylsulfonyl (1-6C)alkyl, sulfamoyl (1-6C)alkyl, N-(1-6C)alkyl-sulfamoyl (1-6C)alkyl, N,N-di-[(1-6C)alkyl]sulfamoyl (1-6C)alkyl, (2-6C)alkanoyl (1-6C)alkyl, (2-6C)alkanoyloxy (1-6C)alkyl or (1-6C)alkoxy-carbonyl (1-6C)alkyl, and

wherein any CH₂ or CH₃ group within -X¹-Q¹ optionally bears on each said CH₂ or CH₃ group one or more halogeno or (1-6C)alkyl substituents or a substituent selected from hydroxy, cyano, amino, (1-4C)alkoxy, (1-4C)alkylamino and di-[(1-4C)alkylamino];

R⁴, R^{4a}, R⁵ and R^{5a}, which may be the same or different, are selected from hydrogen and (1-6C)alkyl, or

R^4 and R^{4a} together with the carbon atom to which they are attached form a (3-7C)cycloalkyl ring, or

R^5 and R^{5a} together with the carbon atom to which they are attached form a (3-7C)cycloalkyl ring, and

wherein any CH_2 or CH_3 group within any of R^4 , R^{4a} , R^5 and R^{5a} optionally bears on each said

CH_2 or CH_3 group one or more halogeno substituents or a substituent selected from hydroxy, cyano, (1-6C)alkoxy, amino, (2-6C)alkanoyl, (1-6C)alkylamino and di-[(1-6C)alkylamino];

R^6 is selected from hydrogen, (1-6C)alkyl, (2-6C)alkenyl, (2-6C)alkynyl, (3-7C)cycloalkyl, (3-7C)cycloalkyl-(1-6C)alkyl, (3-7C)cycloalkenyl, (3-7C)cycloalkenyl-(1-6C)alkyl, heterocyclyl and heterocyclyl-(1-6C)alkyl, and

wherein any heterocyclyl group within an R^6 substituent optionally bears one or more

substituents, which may be the same or different, selected from halogeno, trifluoromethyl, cyano, nitro, hydroxy, amino, formyl, mercapto, (1-6C)alkyl, (2-6C)alkenyl, (2-6C)alkynyl, (1-6C)alkoxy, (1-6C)alkylthio, (1-6C)alkylsulfinyl, (1-6C)alkylsulfonyl, (1-6C)alkylamino, di-[(1-6C)alkyl]amino, (2-6C)alkanoyl, (2-6C)alkanoyloxy and from a group of the formula:

X^3-R^{10} , wherein

X^3 is a direct bond or is selected from O, CO, SO_2 and $N(R^{11})$, wherein;

R^{11} is hydrogen or (1-4C)alkyl, and;

R^{10} is halogeno-(1-4C)alkyl, hydroxy-(1-4C)alkyl, (1-4C)alkoxy-(1-4C)alkyl, cyano-(1-4C)alkyl, amino-(1-4C)alkyl, N-(1-4C)alkylamino-(1-4C)alkyl and N,N-di-[(1-4C)alkyl]amino-(1-4C)alkyl, and

wherein any heterocyclyl group within an R^6 substituent optionally bears 1 or 2 oxo or thioxo substituents, and

wherein any CH_2 or CH_3 group within a R^6 substituent, other than a CH_2 group within a heterocyclyl group, optionally bears on each said CH_2 or CH_3 group one or more halogeno or (1-6C)alkyl substituents or a substituent selected from hydroxy, cyano, amino, carboxy, carbamoyl, sulfamoyl, (2-6C)alkenyl, (2-6C)alkynyl, (1-6C)alkoxy, (1-6C)alkylthio, (1-6C)alkylsulfinyl, (1-6C)alkylsulfonyl, (1-6C)alkylamino, di-[(1-6C)alkyl]amino, N-(1-6C)alkylcarbamoyl, N,N-di-[(1-6C)alkyl]carbamoyl, (2-6C)alkanoyl,

(2-6C)alkanoyloxy, (2-6C)alkanoylamino, $\underline{\text{N}}$ -(1-6C)alkyl-(2-6C)alkanoylamino, $\underline{\text{N}}$ -(1-6C)alkylsulfamoyl, $\underline{\text{N}}$, $\underline{\text{N}}$ -di-[(1-6C)alkyl]sulfamoyl, (1-6C)alkanesulfonylamino and $\underline{\text{N}}$ -(1-6C)alkyl-(1-6C)alkanesulfonylamino;

A is selected from hydrogen, a group of the formula $\text{Z}-(\text{CR}^{12}\text{R}^{13})_p$ and R^{14}_{τ} ; wherein p is 1, 2, 3, or 4;

each R^{12} and R^{13} , which may be the same or different, is selected from hydrogen, (1-6C)alkyl, (2-6C)alkenyl and (2-6C)alkynyl, or an R^{12} and an R^{13} group attached to the same carbon atom form a (3-7C)cycloalkyl or (3-7C)cycloalkenyl ring, and

wherein any CH_2 or CH_3 group within any of R^{12} and R^{13} , optionally bears on each said CH_2 or CH_3 group one or more halogeno or (1-6C)alkyl substituents or a substituent selected from hydroxy, cyano, (1-6C)alkyl, (1-6C)alkoxy, amino, (2-6C)alkanoyl, (1-6C)alkylamino and di-[(1-6C)alkyl]amino;

Z is selected from hydrogen, OR^{15} , $\text{NR}^{16}\text{R}^{17}$, (1-6C)alkylsulfonyl, (1-6C)alkanesulfonylamino and $\underline{\text{N}}$ -(1-6C)alkyl-(1-6C)alkanesulfonylamino; wherein;

each of R^{15} , R^{16} and R^{17} , which may be the same or different, is selected from hydrogen, (1-6C)alkyl, (2-6C)alkenyl, (2-6C)alkynyl and (1-6C)alkoxycarbonyl,

or Z is a group of the formula: $\text{Q}^2\text{-X}^4$ - wherein;

X^4 is selected from O, $\text{N}(\text{R}^{18})$, SO_2 and $\text{SO}_2\text{N}(\text{R}^{18})_{\tau}$; wherein;

R^{18} is hydrogen or (1-6C)alkyl; and;

Q^2 is (3-7C)cycloalkyl, (3-7C)cycloalkenyl or heterocyclyl;

R^{14} is selected from hydrogen, OR^{19} and $\text{NR}^{16}\text{R}^{17}$; wherein;

R^{19} is selected from (1-6C)alkyl, (2-6C)alkenyl and (2-6C)alkynyl, and wherein R^{16} and R^{17} are as defined above,

or R^{14} is a group of the formula: $\text{Q}^3\text{-X}^5$ - wherein;

X^5 is selected from O and $\text{N}(\text{R}^{20})_{\tau}$; wherein;

R^{20} is hydrogen or (1-6C)alkyl; and;

Q^3 is (3-7C)cycloalkyl, (3-7C)cycloalkyl-(1-6C)alkyl, (3-7C)cycloalkenyl, (3-7C)cycloalkenyl-(1-6C)alkyl, heterocyclyl and heterocyclyl-(1-6C)alkyl,

or R^{14} is Q^4 wherein:

Q^4 is (3-7C)cycloalkyl, (3-7C)cycloalkyl-(1-6C)alkyl, (3-7C)cycloalkenyl, (3-7C)cycloalkenyl-(1-6C)alkyl, heterocyclyl or heterocyclyl-(1-6C)alkyl, and

wherein adjacent carbon atoms in any (2-6C)alkylene chain within a Z or R^{14} substituent are optionally separated by the insertion into the chain of a group selected from O, S, SO, SO_2 ,

$N(R^{21})$, CO, -C=C- and -C \equiv C-, wherein:

R^{21} is hydrogen or (1-6C)alkyl, and

wherein any heterocyclyl group within a Z or R^{14} substituent optionally bears one or more substituents, which may be the same or different, selected from halogeno, trifluoromethyl, cyano, nitro, hydroxy, amino, formyl, mercapto, (1-6C)alkyl, (2-6C)alkenyl, (2-6C)alkynyl, (1-6C)alkoxy, (1-6C)alkylthio, (1-6C)alkylsulfanyl, (1-6C)alkylsulfonyl, (1-6C)alkylamino, di-[(1-6C)alkyl]amino, (2-6C)alkanoyl, (2-6C)alkanoyloxy and from a group of the formula:

- X^6 - R^{22} , wherein

X^6 is a direct bond or is selected from O, CO, SO_2 and $N(R^{23})$, wherein:

R^{23} is hydrogen or (1-4C)alkyl, and:

R^{22} is halogeno-(1-4C)alkyl, hydroxy-(1-4C)alkyl, (1-4C)alkoxy-(1-4C)alkyl, cyano-(1-4C)alkyl, amino-(1-4C)alkyl, N-(1-4C)alkylamino-(1-4C)alkyl and N,N-di-[(1-4C)alkyl]amino-(1-4C)alkyl, and

wherein any heterocyclyl group within a Z or R^{14} substituent optionally bears 1 or 2 oxo or thioxo substituents, and

wherein any CH_2 or CH_3 group within a Z or R^{14} group, other than a CH_2 group within a heterocyclyl ring, optionally bears on each said CH_2 or CH_3 group one or more halogeno or (1-6C)alkyl substituents or a substituent selected from hydroxy, cyano, amino, carboxy, carbamoyl, sulfamoyl, (2-6C)alkenyl, (2-6C)alkynyl, (1-6C)alkoxy, (1-6C)alkylthio, (1-6C)alkylsulfanyl, (1-6C)alkylsulfonyl, (1-6C)alkylamino, di-[(1-6C)alkyl]amino, N-(1-6C)alkylcarbamoyl, N,N-di-[(1-6C)alkyl]carbamoyl, (2-6C)alkanoyl, (2-6C)alkanoyloxy, (2-6C)alkanoylamino, N-(1-6C)alkyl-(2-6C)alkanoylamino, N-(1-6C)alkylsulfamoyl, N,N-di-[(1-6C)alkyl]sulfamoyl, (1-6C)alkanesulfonylamino and N-(1-6C)alkyl-(1-6C)alkanesulfonylamino;

or a pharmaceutically acceptable salt thereof.

Claim 2 (currently amended): A ~~The~~ quinazoline derivative according to claim 1, wherein:
 m is 0, 1 or 2;

each R^1 , which may be the same or different, is selected from hydroxy, (1-6C)alkoxy,

(3-7C)cycloalkyl-oxy and (3-7C)cycloalkyl-(1-6C)alkoxy, and

wherein any CH_2 or CH_3 group within a R^1 substituent optionally bears on each said CH_2 or CH_3 group one or more halogeno or (1-6C)alkyl substituents, or a substituent selected from hydroxy and (1-6C)alkoxy,

R^2 is hydrogen or (1-4C)alkyl;

n is 0, 1, 2, 3 or 4;

each R^3 , which may be the same or different, is selected from halogeno, (1-4C)alkyl, trifluoromethyl, (1-4C)alkoxy, (2-4C)alkenyl and (2-4C)alkynyl;

X^1 is selected from O, S, SO, SO_2 , $N(R^7)$, $CH(OR^7)$, $CON(R^7)$, $N(R^7)CO$, $SO_2N(R^7)$, $N(R^7)SO_2$, $OC(R^7)_2$, $C(R^7)_2O$, $SC(R^7)_2$, $C(R^7)_2S$, CO, $C(R^7)_2N(R^7)$ and $N(R^7)C(R^7)_2$, wherein;

each R^7 , which may be the same or different, is hydrogen or (1-6C)alkyl;

Q^1 is aryl, or heteroaryl, and

wherein Q^1 optionally bears one or more substituents, which may be the same or different, selected from halogeno, cyano, nitro, hydroxy, amino, carboxy, carbamoyl, sulfamoyl, formyl, mercapto, (1-6C)(1-4C)alkyl, (2-8C)alkenyl, (2-8C)alkynyl, and (1-6C)(1-4C)alkoxy,

(2-6C)alkenyl-oxy, (2-6C)alkynyl-oxy, (1-6C)alkylthio, (1-6C)alkylsulfanyl,

(1-6C)alkylsulfonyl, (1-6C)alkylamino, di-[(1-6C)alkyl]amino, (1-6C)alkoxy-carbonyl,

N -(1-6C)alkyl-carbamoyl, N,N -di-[(1-6C)alkyl]carbamoyl, (2-6C)alkanoyl, (3-6C)alkenoyl,

(3-6C)alkynoyl, (2-6C)alkanoyl-oxy, (2-6C)alkanoylamino,

N -(1-6C)alkyl-(2-6C)alkanoylamino, (3-6C)alkenoylamino, N -(1-6C)alkyl-(3-

6C)alkenoylamino, (3-6C)alkynoylamino, N -(1-6C)alkyl-(3-6C)alkynoylamino,

N -(1-6C)alkyl-sulfamoyl, N,N -di-[(1-6C)alkyl]sulfamoyl, (1-6C)alkanesulfonylamino,

N -(1-6C)alkyl-(1-6C)alkanesulfonylamino, and a group of the formula: $-X^2-R^8$ wherein

X^2 is a direct bond or is selected from O, CO and $N(R^9)$, wherein

R^9 is hydrogen or (1-6C)alkyl, and

R^8 is halogeno (1-6C)alkyl, hydroxy (1-6C)alkyl, carboxy (1-6C)alkyl, (1-6C)alkoxy (1-6C)alkyl, cyano (1-6C)alkyl, amino (1-6C)alkyl, N-(1-6C)alkylamino (1-6C)alkyl, N,N-di-[(1-6C)alkyl]amino (1-6C)alkyl, (2-6C)alkanoylamino (1-6C)alkyl, N-(1-6C)alkyl (2-6C)alkanoylamino (1-6C)alkyl, (1-6C)alkoxy-carbonylamino (1-6C)alkyl, carbamoyl (1-6C)alkyl, N-(1-6C)alkyl-carbamoyl (1-6C)alkyl, N,N-di-[(1-6C)alkyl]carbamoyl (1-6C)alkyl, (1-6C)alkylthio (1-6C)alkyl, (1-6C)alkylsulfinyl (1-6C)alkyl, (1-6C)alkylsulfonyl (1-6C)alkyl, sulfamoyl (1-6C)alkyl, N-(1-6C)alkylsulfamoyl (1-6C)alkyl, N,N-di-[(1-6C)alkylsulfamoyl (1-6C)alkyl, (2-6C)alkanoyl (1-6C)alkyl, (2-6C)alkanoyloxy (1-6C)alkyl or (1-6C)alkoxy-carbonyl (1-6C)alkyl, and

wherein any CH_2 or CH_3 group within $-X^1-Q^1$ optionally bears on each said CH_2 or CH_3 group one or more halogeno or (1-6C)alkyl substituents or a substituent selected from hydroxy, cyano, amino, (1-4C)alkoxy, (1-4C)alkylamino and di-[(1-4C)alkylamino];

R^4 , R^{4a} , R^5 and R^{5a} , which may be the same or different, are selected from hydrogen and (1-6C)alkyl, or

R^4 and R^{4a} together with the carbon atom to which they are attached form a (3-7C)cycloalkyl ring, or

R^5 and R^{5a} together with the carbon atom to which they are attached form a (3-7C)cycloalkyl ring, and

wherein any CH_2 or CH_3 group within any of R^4 , R^{4a} , R^5 and R^{5a} optionally bears on each said CH_2 or CH_3 group one or more halogeno substituents or a substituent selected from hydroxy, cyano, (1-6C)alkoxy, amino, (2-6C)alkanoyl, (1-6C)alkylamino and di-[(1-6C)alkylamino];

R^6 is selected from hydrogen, (1-6C)alkyl, (2-6C)alkenyl, (2-6C)alkynyl, (3-7C)cycloalkyl, (3-7C)cycloalkyl-(1-6C)alkyl, (3-7C)cycloalkenyl, (3-7C)cycloalkenyl-(1-6C)alkyl, heterocyclyl and heterocyclyl-(1-6C)alkyl, and

wherein any heterocyclyl group within an R^6 substituent optionally bears one or more substituents, which may be the same or different, selected from halogeno, trifluoromethyl, cyano, nitro, hydroxy, amino, formyl, mercapto, (1-6C)alkyl, (2-6C)alkenyl, (2-6C)alkynyl,

(1-6C)alkoxy, (1-6C)alkylthio, (1-6C)alkylsulfanyl, (1-6C)alkylsulfonyl, (1-6C)alkylamino, di-[(1-6C)alkyl]amino, (2-6C)alkanoyl, (2-6C)alkanoyloxy and from a group of the formula:
 $-X^3-R^{10}$; wherein

X^3 is a direct bond or is selected from O, CO, SO₂ and N(R¹¹), wherein:

R¹¹ is hydrogen or (1-4C)alkyl, and R¹⁰ is halogeno-(1-4C)alkyl, hydroxy-(1-4C)alkyl, (1-4C)alkoxy-(1-4C)alkyl, cyano-(1-4C)alkyl, amino-(1-4C)alkyl,

N-(1-4C)alkylamino-(1-4C)alkyl and N,N-di-[(1-4C)alkyl]amino-(1-4C)alkyl, and

wherein any heterocyclyl group within an R⁶ substituent optionally bears 1 or 2 oxo or thioxo substituents; and

wherein any CH₂ or CH₃ group within a R⁶ substituent, other than a CH₂ group within a heterocyclyl group, optionally bears on each said CH₂ or CH₃ group one or more halogeno or (1-6C)alkyl substituents or a substituent selected from hydroxy, cyano, amino, carboxy, carbamoyl, sulfamoyl, (2-6C)alkenyl, (2-6C)alkynyl, (1-6C)alkoxy, (1-6C)alkylthio, (1-6C)alkylsulfanyl, (1-6C)alkylsulfonyl, (1-6C)alkylamino, di-[(1-6C)alkyl]amino, N-(1-6C)alkylcarbamoyl, N,N-di-[(1-6C)alkyl]carbamoyl, (2-6C)alkanoyl, (2-6C)alkanoyloxy, (2-6C)alkanoylamino, N-(1-6C)alkyl-(2-6C)alkanoylamino, N-(1-6C)alkylsulfamoyl, N,N-di-[(1-6C)alkyl]sulfamoyl, (1-6C)alkanesulfonylamino and N-(1-6C)alkyl-(1-6C)alkanesulfonylamino;

A is selected from hydrogen, a group of the formula Z-(CR¹²R¹³)_p- and R¹⁴; wherein

p is 1, 2, 3, or 4,

each R¹² and R¹³, which may be the same or different, is selected from hydrogen, (1-6C)alkyl, (2-6C)alkenyl and (2-6C)alkynyl, or an R¹² and an R¹³ group attached to the same carbon atom form a (3-7C)cycloalkyl or (3-7C)cycloalkenyl ring, and

wherein any CH₂ or CH₃ group within any of R¹² and R¹³, optionally bears on each said CH₂ or CH₃ group one or more halogeno or (1-6C)alkyl substituents or a substituent selected from hydroxy, cyano, (1-6C)alkyl, (1-6C)alkoxy, amino, (2-6C)alkanoyl, (1-6C)alkylamino and di-[(1-6C)alkyl]amino;

Z is selected from hydrogen, OR¹⁵, NR¹⁶R¹⁷, (1-6C)alkylsulfonyl, (1-6C)alkanesulfonylamino

and $\text{N}-(1-6\text{C})\text{alkyl}-(1-6\text{C})\text{alkanesulfonylamino}$, wherein;

each of R¹⁵, R¹⁶ and R¹⁷, which may be the same or different, is selected from hydrogen, (1-6C)alkyl, (2-6C)alkenyl and (2-6C)alkynyl,

or Z is a group of the formula: Q²-X⁴-, wherein

X⁴ is selected from O, N(R¹⁸), SO₂ and SO₂N(R¹⁸), wherein;

R¹⁸ is hydrogen or (1-6C)alkyl, and;

Q² is (3-7C)cycloalkyl, (3-7C)cycloalkenyl or heterocyclyl;

R¹⁴ is selected from hydrogen, OR¹⁹ and NR¹⁶R¹⁷, wherein;

R¹⁹ is selected from (1-6C)alkyl, (2-6C)alkenyl and (2-6C)alkynyl, and wherein R¹⁶ and R¹⁷ are as defined above,

or R¹⁴ is a group of the formula: Q³-X⁵-, wherein

X⁵ is selected from O and N(R²⁰), wherein R²⁰ is hydrogen or (1-6C)alkyl, and;

Q³ is (3-7C)cycloalkyl, (3-7C)cycloalkyl-(1-6C)alkyl, (3-7C)cycloalkenyl, (3-7C)cycloalkenyl-(1-6C)alkyl, heterocyclyl and heterocyclyl-(1-6C)alkyl,

or R¹⁴ is Q⁴ wherein Q⁴ is (3-7C)cycloalkyl, (3-7C)cycloalkenyl or heterocyclyl, and

wherein adjacent carbon atoms in any (2-6C)alkylene chain within a Z or R¹⁴ substituent are optionally separated by the insertion into the chain of a group selected from O, S, SO, SO₂,

N(R²¹), CO, -C=C- and -C≡C-, wherein;

R²¹ is hydrogen or (1-6C)alkyl, and

wherein any heterocyclyl group within a Z or R¹⁴ substituent optionally bears one or more

substituents, which may be the same or different, selected from halogeno, trifluoromethyl,

cyano, nitro, hydroxy, amino, formyl, mercapto, (1-6C)alkyl, (2-6C)alkenyl, (2-6C)alkynyl,

(1-6C)alkoxy, (1-6C)alkylthio, (1-6C)alkylsulfinyl, (1-6C)alkylsulfonyl, (1-6C)alkylamino,

di-[(1-6C)alkyl]amino, (2-6C)alkanoyl, (2-6C)alkanoyloxy and from a group of the formula:

-X⁶-R²²-, wherein

X⁶ is a direct bond or is selected from O, CO, SO₂ and N(R²³), wherein;

R²³ is hydrogen or (1-4C)alkyl, and;

R^{22} is halogeno-(1-4C)alkyl, hydroxy-(1-4C)alkyl, (1-4C)alkoxy-(1-4C)alkyl, cyano-(1-4C)alkyl, amino-(1-4C)alkyl, N -(1-4C)alkylamino-(1-4C)alkyl and N,N -di-[(1-4C)alkyl]amino-(1-4C)alkyl, and wherein any heterocyclyl group within a Z or R^{14} substituent optionally bears 1 or 2 oxo or thioxo substituents, and wherein any CH_2 or CH_3 group within a Z or R^{14} group, other than a CH_2 group within a heterocyclyl ring, optionally bears on each said CH_2 or CH_3 group one or more halogeno or (1-6C)alkyl substituents or a substituent selected from hydroxy, cyano, amino, carboxy, carbamoyl, sulfamoyl, (2-6C)alkenyl, (2-6C)alkynyl, (1-6C)alkoxy, (1-6C)alkylthio, (1-6C)alkylsulfanyl, (1-6C)alkylsulfonyl, (1-6C)alkylamino, di-[(1-6C)alkyl]amino, N -(1-6C)alkylcarbamoyl, N,N -di-[(1-6C)alkyl]carbamoyl, (2-6C)alkanoyl, (2-6C)alkanoyloxy, (2-6C)alkanoylamino, N -(1-6C)alkyl-(2-6C)alkanoylamino, N -(1-6C)alkylsulfamoyl, N,N -di-[(1-6C)alkyl]sulfamoyl, (1-6C)alkanesulfonylamino and N -(1-6C)alkyl-(1-6C)alkanesulfonylamino; or a pharmaceutically acceptable salt thereof.

Claim 3 (currently amended): A The quinazoline derivative according to claim 1, wherein R^4 , R^{4a} , R^5 and R^{5a} , which may be the same or different, are selected from hydrogen and (1-6C)alkyl, and wherein any CH_2 or CH_3 group within any of R^4 , R^{4a} , R^5 and R^{5a} optionally bears on each said CH_2 or CH_3 group one or more halogeno substituents or a substituent selected from hydroxy, cyano, (1-6C)alkoxy, amino, (2-6C)alkanoyl, (1-6C)alkylamino and di-[(1-6C)alkylamino].

Claim 4 (currently amended): A The quinazoline derivative according to claim 1, wherein m is 0.

Claim 5 (currently amended): A The quinazoline derivative according to claim 1, wherein R^2 is hydrogen.

Claim 6 (currently amended): ~~A-~~The quinazoline derivative according to claim 1, wherein n is 0, 1 or 2 and, when present, at least one R³ is in a meta-position (3-position) relative to the nitrogen of the anilino group in formula I.

Claim 7 (currently amended): ~~A-~~The quinazoline derivative according to claim 1, wherein n is 1 and R³ is selected from halogeno and (1-4C)alkyl.

Claim 8 (currently amended): ~~A-~~The quinazoline derivative according to claim 7, wherein R³ is chloro.

Claim 9 (currently amended): ~~A-~~The quinazoline derivative according to claim 7, wherein R³ is methyl.

Claim 10 (currently amended): ~~A-~~The quinazoline derivative according to claim 1, wherein X¹ is selected from O, S, OC(R⁷)₂, SC(R⁷)₂, SO, SO₂, N(R⁷), CO and N(R⁷)C(R⁷)₂ wherein each R⁷, which may be the same or different, is selected from hydrogen or (1-6C)alkyl.

Claim 11 (currently amended): ~~A-~~The quinazoline derivative according to claim 1, wherein X¹ is selected from O, S and OC(R⁷)₂ wherein each R⁷ is, independently, hydrogen or (1-4C)alkyl.

Claim 12 (currently amended): ~~A-~~The quinazoline derivative according to claim 1, wherein X¹ is OCH₂.

Claim 13 (currently amended): ~~A-~~The quinazoline derivative according to claim 1, wherein

Q¹ is selected from phenyl and a 5- or 6-membered monocyclic heteroaryl ring, which ring contains 1, 2 or 3 heteroatoms independently selected from oxygen, nitrogen and sulfur, and wherein Q¹ optionally bears one or more substituents, which may be the same or different, selected from halogeno, cyano, nitro, hydroxy, amino, carboxy, carbamoyl, sulfamoyl, formyl,

mercapto, (1-6C)(1-4C)alkyl, (2-8C)alkenyl, (2-8C)alkynyl, and (1-6C)(1-4C)alkoxy, (2-6C)alkenyloxy, (2-6C)alkynyloxy, (1-6C)alkylthio, (1-6C)alkylsulfanyl, (1-6C)alkylsulfonyl, (1-6C)alkylamino, di-[(1-6C)alkyl]amino, (1-6C)alkoxy-carbonyl, N-(1-6C)alkyl-carbamoyl, N,N-di-[(1-6C)alkyl]carbamoyl, (2-6C)alkanoyl, (3-6C)alkenoyl, (3-6C)alkynoyl, (2-6C)alkanoyloxy, (2-6C)alkanoylamino, N-(1-6C)alkyl-(2-6C)alkanoylamino, (3-6C)alkenoylamino, N-(1-6C)alkyl-(3-6C)alkenoylamino, (3-6C)alkynoylamino, N-(1-6C)alkyl-(3-6C)alkynoylamino, N-(1-6C)alkylsulfamoyl, N,N-di-[(1-6C)alkyl]sulfamoyl, (1-6C)alkanesulfonylamino, N-(1-6C)alkyl-(1-6C)alkanesulfonylamino, and a group of the formula: $-X^2-R^8$ wherein X^2 is a direct bond or is selected from O, CO and N(R^9), wherein R^9 is hydrogen or (1-6C)alkyl, and R^8 is halogeno (1-6C)alkyl, hydroxy (1-6C)alkyl, carboxy (1-6C)alkyl, (1-6C)alkoxy (1-6C)alkyl, cyano (1-6C)alkyl, amino (1-6C)alkyl, N-(1-6C)alkylamino (1-6C)alkyl, N,N-di-[(1-6C)alkyl]amino (1-6C)alkyl, (2-6C)alkanoylamino (1-6C)alkyl, N-(1-6C)alkyl-(2-6C)alkanoylamino (1-6C)alkyl, (1-6C)alkoxy-carbonylamino (1-6C)alkyl, carbamoyl (1-6C)alkyl, N-(1-6C)alkyl-carbamoyl (1-6C)alkyl, N,N-di-[(1-6C)alkyl]carbamoyl (1-6C)alkyl, (1-6C)alkylthio (1-6C)alkyl, (1-6C)alkylsulfanyl (1-6C)alkyl, (1-6C)alkylsulfonyl (1-6C)alkyl, sulfamoyl (1-6C)alkyl, N-(1-6C)alkylsulfamoyl (1-6C)alkyl, N,N-di-[(1-6C)alkyl]sulfamoyl (1-6C)alkyl, (2-6C)alkanoyl (1-6C)alkyl, (2-6C)alkanoyloxy (1-6C)alkyl or (1-6C)alkoxy-carbonyl (1-6C)alkyl, and

wherein any CH_2 or CH_3 group within $-X^1-Q^1$ optionally bears on each said CH_2 or CH_3 group one or more halogeno or (1-6C)alkyl substituents or a substituent selected from hydroxy, cyano, amino, (1-4C)alkoxy, (1-4C)alkylamino and di-[(1-4C)alkylamino].

Claim 14 (currently amended): A The quinazoline derivative according to claim 1, wherein Q^1 is selected from phenyl, pyridyl, pyrazinyl, 1,3-thiazolyl, 1H-imidazolyl, 1H-pyrazolyl, 1,3-oxazolyl and isoxazolyl.

Claim 15 (currently amended): ~~A~~The quinazoline derivative according to claim 1,
wherein

R^6 is selected from hydrogen, (1-3C)alkyl, (2-3C)alkenyl, (2-3C)alkynyl, (3-5C)cycloalkyl, (3-5C)cycloalkyl-(1-3C)alkyl, heterocyclyl and heterocyclyl-(1-3C)alkyl,

wherein any heterocyclyl group within R^6 is a 4, 5, 6 or 7 membered monocyclic saturated or partially saturated heterocyclyl ring containing 1 or 2 heteroatoms selected from oxygen, nitrogen and sulfur, which heterocyclyl group is linked to the group to which it is attached by a ring carbon atom, and

wherein any heterocyclyl group within an R^6 substituent optionally bears one or more substituents, which may be the same or different, selected from halogeno, trifluoromethyl, cyano, nitro, hydroxy, amino, mercapto, (1-6C)alkyl, (2-6C)alkenyl, (2-6C)alkynyl, (1-6C)alkoxy, (1-6C)alkylthio, (1-6C)alkylsulfinyl, (1-6C)alkylsulfonyl, (1-6C)alkylamino, di-[(1-6C)alkyl]amino, (2-6C)alkanoyl, (2-6C)alkanoyloxy and from a group of the formula:

$-X^3-R^{10}$; ~~wherein~~

X^3 is a direct bond or is selected from O and N(R^{11}); ~~wherein;~~

R^{11} is hydrogen or (1-4C)alkyl; ~~and;~~

R^{10} is halogeno-(1-4C)alkyl, hydroxy-(1-4C)alkyl, (1-4C)alkoxy-(1-4C)alkyl, cyano-(1-4C)alkyl, amino-(1-4C)alkyl, N-(1-4C)alkylamino-(1-4C)alkyl and N,N-di-[(1-4C)alkyl]amino-(1-4C)alkyl, ~~and~~

wherein any heterocyclyl group within an R^6 substituent optionally bears 1 or 2 oxo substituents; and wherein any CH_2 or CH_3 group within a R^6 substituent, other than a CH_2 group within a heterocyclyl group, optionally bears on each said CH_2 or CH_3 group one or more halogeno or (1-6C)alkyl substituents or a substituent selected from hydroxy, amino, (1-6C)alkoxy, (1-6C)alkylamino and di-[(1-6C)alkyl]amino.

Claim 16 (currently amended): ~~A~~The quinazoline derivative according to claim 15, wherein R^6 is (1-3C)alkyl, and wherein any CH_2 or CH_3 group within a R^6 substituent, other than a CH_2 group within a heterocyclyl group, optionally bears on each said CH_2 or CH_3 group one or more

halogeno or (1-6C)alkyl substituents or a substituent selected from hydroxy, amino, (1-6C)alkoxy, (1-6C)alkylamino and di-[(1-6C)alkyl]amino.

Claim 17 (currently amended): ~~A~~The quinazoline derivative according to claim 1, wherein

A is selected from a group of the formula $Z-(CR^{12}R^{13})_p-$ and R^{14} , wherein

p is 1, 2 or 3;

each R^{12} and R^{13} , which may be the same or different, is selected from hydrogen and

(1-6C)alkyl, and

wherein any CH_2 or CH_3 group within any of R^{12} and R^{13} optionally bears on each said CH_2 or CH_3 group one or more halogeno substituents or a substituent selected from hydroxy and (1-6C)alkoxy;

Z is selected from hydrogen, OR^{15} , $NR^{16}R^{17}$ and (1-6C)alkylsulfonyl, wherein;

each of R^{15} , R^{16} and R^{17} , which may be the same or different, is selected from hydrogen, (1-6C)alkyl and (1-6C)alkoxycarbonyl;

R^{14} is selected from OR^{19} and $NR^{16}R^{17}$, wherein;

R^{19} is selected from (1-6C)alkyl and wherein R^{16} and R^{17} are as defined above,

or R^{14} is Q^4 wherein;

Q^4 is (3-7C)cycloalkyl, heterocyclcyl or heterocyclcyl-(1-6C)alkyl, and

wherein any heterocyclcyl group within a Z or R^{14} substituent optionally bears one or more substituents, which may be the same or different, selected from halogeno, hydroxy, (1-6C)alkyl and (1-6C)alkoxy, and

wherein any CH_2 or CH_3 group within a Z or R^{14} group, other than a CH_2 group within a heterocyclcyl ring, optionally bears on each said CH_2 or CH_3 group one or more halogeno or (1-6C)alkyl substituents or a substituent selected from hydroxy and (1-6C)alkoxy.

Claim 18 (currently amended): ~~A~~The quinazoline derivative selected from one or more of the following:

- N*-{2-[(4-{3-chloro-4-(pyridin-2-ylmethoxy)anilino}quinazolin-5-yl)oxy]ethyl}-2-methoxy-*N*-methylacetamide;
- N*-{2-[(4-{3-chloro-4-(pyridin-2-ylmethoxy)anilino}quinazolin-5-yl)oxy]ethyl}-2-(dimethylamino)-*N*-methylacetamide;
- N*-{(2*R*)-2-[(4-{3-chloro-4-(pyridin-2-ylmethoxy)anilino}quinazolin-5-yl)oxy]propyl}-2-methoxy-*N*-methylacetamide;
- 2-hydroxy-*N*-methyl-*N*-{2-[(4-{3-methyl-4-(pyrazin-2-ylmethoxy)anilino}quinazolin-5-yl)oxy]ethyl}acetamide;
- 2-hydroxy-*N*-methyl-*N*-{2-[(4-{3-methyl-4-(1,3-thiazol-4-ylmethoxy)anilino}quinazolin-5-yl)oxy]ethyl}acetamide;
- 2-hydroxy-*N*-methyl-*N*-(2-[(4-{3-methyl-4-[(5-methylisoxazol-3-yl)methoxy]anilino}quinazolin-5-yl)oxy]ethyl)acetamide;
- N*-{(2*R*)-2-[(4-{3-chloro-4-(pyridin-2-ylmethoxy)anilino}quinazolin-5-yl)oxy]propyl}-2-methoxyacetamide;
- N*-(2-[[4-(3-chloro-4-[(6-methylpyridin-2-yl)methoxy]anilino)quinazolin-5-yl]oxy]ethyl)-2-hydroxy-*N*-methylacetamide;
- N*-((2*R*)-2-[[4-(3-chloro-4-[(6-methylpyridin-2-yl)methoxy]anilino)quinazolin-5-yl]oxy]propyl)-2-hydroxy-*N*-methylacetamide;
- N*-(2-[[4-(3-chloro-4-[(6-methylpyridin-2-yl)methoxy]anilino)quinazolin-5-yl]oxy]ethyl)-*N*-methylacetamide;
- N*-(2-[[4-(3-chloro-4-[(2-fluorobenzyl)oxy]anilino)quinazolin-5-yl]oxy]ethyl)-*N*-methylacetamide;
- N*-(2-[[4-(3-chloro-4-[(3-fluorobenzyl)oxy]anilino)quinazolin-5-yl]oxy]ethyl)-*N*-methylacetamide;
- N*-{2-[(4-{3-chloro-4-(1,3-thiazol-4-ylmethoxy)anilino}quinazolin-5-yl)oxy]ethyl}-*N*-methylacetamide;
- N*-{2-[(4-{3-chloro-4-(pyrazin-2-ylmethoxy)anilino}quinazolin-5-yl)oxy]ethyl}-*N*-methylacetamide;

- N*-{(2*R*)-2-[(4-{3-chloro-4-(pyridin-2-ylmethoxy)anilino}quinazolin-5-yl)oxy]propyl}-2-hydroxyacetamide;
- N*-{2-[(4-{3-chloro-4-(pyridin-2-ylmethoxy)anilino}quinazolin-5-yl)oxy]ethyl}-*N*-methylacetamide;
- 2-hydroxy-*N*-methyl-*N*-{2-[(4-{3-methyl-4-(pyridin-2-ylmethoxy)anilino}quinazolin-5-yl)oxy]ethyl}acetamide;
- N*-{(1*R*)-2-[(4-{3-chloro-4-(pyridin-2-ylmethoxy)anilino}quinazolin-5-yl)oxy]-1-methylethyl}acetamide;
- N*-{(1*R*)-2-[(4-{3-chloro-4-(pyridin-2-ylmethoxy)anilino}quinazolin-5-yl)oxy]-1-methylethyl}-2-hydroxyacetamide;
- N*-{2-[(4-{3-chloro-4-(pyridin-2-ylmethoxy)anilino}quinazolin-5-yl)oxy]ethyl}-2-hydroxy-*N*-methylacetamide;
- N*-(2-{[4-(3-chloro-4-[(3-fluorobenzyl)oxy]anilino)quinazolin-5-yl]oxy}ethyl)-2-hydroxy-*N*-methylacetamide;
- N*-{2-[(4-{3-chloro-4-(1,3-thiazol-4-ylmethoxy)anilino}quinazolin-5-yl)oxy]ethyl}-2-hydroxy-*N*-methylacetamide;
- N*-{2-[(4-{3-chloro-4-(pyrazin-2-ylmethoxy)anilino}quinazolin-5-yl)oxy]ethyl}-2-hydroxy-*N*-methylacetamide;
- N*-{2-[(4-{3-chloro-4-(pyridin-2-ylmethoxy)anilino}quinazolin-5-yl)oxy]ethyl}acetamide;
- N*-{(2*R*)-2-[(4-{3-chloro-4-(pyridin-2-ylmethoxy)anilino}quinazolin-5-yl)oxy]propyl}acetamide;
- N*-{(2*R*)-2-[(4-{3-chloro-4-(pyridin-2-ylmethoxy)anilino}quinazolin-5-yl)oxy]propyl}-2-hydroxy-*N*-methylacetamide;
- N*-{(2*R*)-2-[(4-{3-chloro-4-(pyrazin-2-ylmethoxy)anilino}quinazolin-5-yl)oxy]propyl}-2-hydroxy-*N*-methylacetamide;
- N*-((2*R*)-2-{[4-(3-chloro-4-[(3-fluorobenzyl)oxy]anilino)quinazolin-5-yl]oxy}propyl)-2-hydroxy-*N*-methylacetamide;
- N*-{(2*R*)-2-[(4-{3-chloro-4-(1,3-thiazol-4-ylmethoxy)anilino}quinazolin-5-yl)oxy]propyl}-2-hydroxy-*N*-methylacetamide;

N-{(2*R*)-2-[(4-{3-chloro-4-(pyridin-2-ylmethoxy)anilino}quinazolin-5-yl)oxy]propyl}-*N*-methylacetamide;

N-{2-[(4-{[3-chloro-4-(pyridin-2-ylmethoxy)phenyl]amino}quinazolin-5-yl)oxy]ethyl}-*N*-ethylacetamide;

N-{2-[(4-{[3-chloro-4-(pyridin-2-ylmethoxy)phenyl]amino}quinazolin-5-yl)oxy]ethyl}-*N*-ethyl-2-hydroxyacetamide;

N-{2-[(4-{[3-chloro-4-(pyridin-2-ylmethoxy)phenyl]amino}quinazolin-5-yl)oxy]ethyl}-*N*-propylacetamide;

N-{2-[(4-{[3-chloro-4-(pyridin-2-ylmethoxy)phenyl]amino}quinazolin-5-yl)oxy]ethyl}-2-hydroxy-*N*-propylacetamide;

N-{2-[(4-{[3-chloro-4-(pyridin-2-ylmethoxy)phenyl]amino}quinazolin-5-yl)oxy]ethyl}-*N*-isopropylacetamide;

N-{2-[(4-{[3-chloro-4-(pyridin-2-ylmethoxy)phenyl]amino}quinazolin-5-yl)oxy]ethyl}-2-hydroxy-*N*-isopropylacetamide;

N-allyl-*N*-{2-[(4-{[3-chloro-4-(pyridin-2-ylmethoxy)phenyl]amino}quinazolin-5-yl)oxy]ethyl}acetamide;

N-allyl-*N*-{2-[(4-{[3-chloro-4-(pyridin-2-ylmethoxy)phenyl]amino}quinazolin-5-yl)oxy]ethyl}-2-hydroxyacetamide;

N-{2-[(4-{[3-chloro-4-(pyridin-2-ylmethoxy)phenyl]amino}quinazolin-5-yl)oxy]ethyl}-*N*-cyclopropylacetamide;

N-{2-[(4-{[3-chloro-4-(pyridin-2-ylmethoxy)phenyl]amino}quinazolin-5-yl)oxy]ethyl}-*N*-cyclopropyl-2-hydroxyacetamide;

N-{2-[(4-{[3-chloro-4-(pyridin-2-ylmethoxy)phenyl]amino}quinazolin-5-yl)oxy]ethyl}-*N*-(cyclopropylmethyl)acetamide;

N-{2-[(4-{[3-chloro-4-(pyridin-2-ylmethoxy)phenyl]amino}quinazolin-5-yl)oxy]ethyl}-*N*-(cyclopropylmethyl)-2-hydroxyacetamide;

N-{2-[(4-{[3-chloro-4-(pyridin-2-ylmethoxy)phenyl]amino}quinazolin-5-yl)oxy]ethyl}-*N*-cyclobutylacetamide;

- N*-{2-[(4-{[3-chloro-4-(pyridin-2-ylmethoxy)phenyl]amino}quinazolin-5-yl)oxy]ethyl}-*N*-cyclobutyl-2-hydroxyacetamide;
- N*-{2-[(4-{[3-chloro-4-(pyridin-2-ylmethoxy)phenyl]amino}quinazolin-5-yl)oxy]ethyl}-*N*-(1-methylpiperidin-4-yl)acetamide;
- N*-{2-[(4-{[3-chloro-4-(pyridin-2-ylmethoxy)phenyl]amino}quinazolin-5-yl)oxy]ethyl}-*N*-(tetrahydro-2*H*-pyran-4-yl)acetamide;
- N*-{2-[(4-{[3-chloro-4-(pyridin-2-ylmethoxy)phenyl]amino}quinazolin-5-yl)oxy]ethyl}-2-hydroxy-*N*-(tetrahydro-2*H*-pyran-4-yl)acetamide;
- N*-{2-[(4-{[3-chloro-4-(pyridin-2-ylmethoxy)phenyl]amino}quinazolin-5-yl)oxy]ethyl}-*N*-(2-hydroxyethyl)acetamide;
- N*-{2-[(4-{[3-chloro-4-(pyridin-2-ylmethoxy)phenyl]amino}quinazolin-5-yl)oxy]ethyl}-2-hydroxy-*N*-(2-hydroxyethyl)acetamide;
- N*-{2-[(4-{[3-chloro-4-(pyridin-2-ylmethoxy)phenyl]amino}quinazolin-5-yl)oxy]ethyl}-*N*-(2-methoxyethyl)acetamide;
- N*-{2-[(4-{[3-chloro-4-(pyridin-2-ylmethoxy)phenyl]amino}quinazolin-5-yl)oxy]ethyl}-2-hydroxy-*N*-(2-methoxyethyl)acetamide;
- N*-{2-[(4-{[3-chloro-4-(pyridin-2-ylmethoxy)phenyl]amino}quinazolin-5-yl)oxy]ethyl}-*N*-prop-2-yn-1-ylacetamide;
- N*-{2-[(4-{[3-chloro-4-(pyridin-2-ylmethoxy)phenyl]amino}quinazolin-5-yl)oxy]ethyl}-2-hydroxy-*N*-prop-2-yn-1-ylacetamide;
- N*-{2-[(4-{[3-chloro-4-(pyridin-2-ylmethoxy)phenyl]amino}quinazolin-5-yl)oxy]ethyl}-2-hydroxy-*N*-methylpropanamide;
- N*-{2-[(4-{[3-chloro-4-(pyridin-2-ylmethoxy)phenyl]amino}quinazolin-5-yl)oxy]ethyl}-*N*-methyl-tetrahydrofuran-2-carboxamide;
- N*-{2-[(4-{[3-chloro-4-(pyridin-2-ylmethoxy)phenyl]amino}quinazolin-5-yl)oxy]ethyl}-*N*,1-dimethylprolinamide;
- N*-{2-[(4-{[3-chloro-4-(pyridin-2-ylmethoxy)phenyl]amino}quinazolin-5-yl)oxy]ethyl}-2-hydroxy-*N*,2-dimethylpropanamide;

- N*-{2-[(4-{[3-chloro-4-(pyridin-2-ylmethoxy)phenyl]amino}quinazolin-5-yl)oxy]ethyl}-1-hydroxy-*N*-methylcyclopropanecarboxamide;
- N*¹-{2-[(4-{[3-chloro-4-(pyridin-2-ylmethoxy)phenyl]amino}quinazolin-5-yl)oxy]ethyl}-*N*¹,*N*²-dimethylglycinamide;
- N*-{2-[(4-{[3-chloro-4-(pyridin-2-ylmethoxy)phenyl]amino}quinazolin-5-yl)oxy]ethyl}-3-hydroxy-*N*,2,2-trimethylpropanamide;
- N*-{2-[(4-{[3-chloro-4-(pyridin-2-ylmethoxy)phenyl]amino}quinazolin-5-yl)oxy]ethyl}-3-hydroxy-*N*-methylpropanamide;
- N*-{(2*S*)-2-[(4-{[3-chloro-4-(pyridin-2-ylmethoxy)phenyl]amino}quinazolin-5-yl)oxy]propyl}acetamide;
- N*-{(2*S*)-2-[(4-{[3-chloro-4-(pyridin-2-ylmethoxy)phenyl]amino}quinazolin-5-yl)oxy]propyl}-2-hydroxyacetamide;
- N*¹-{(2*S*)-2-[(4-{[3-chloro-4-(pyridin-2-ylmethoxy)phenyl]amino}quinazolin-5-yl)oxy]propyl}-*N*²,*N*²-dimethylglycinamide;
- N*-{(2*S*)-2-[(4-{[3-chloro-4-(pyridin-2-ylmethoxy)phenyl]amino}quinazolin-5-yl)oxy]propyl}-2-methoxyacetamide;
- N*-{(2*S*)-2-[(4-{[3-chloro-4-(pyridin-2-ylmethoxy)phenyl]amino}quinazolin-5-yl)oxy]propyl}-2-(methylsulfonyl)acetamide;
- N*-{2-[(4-{[3-chloro-4-(pyridin-2-ylmethoxy)phenyl]amino}quinazolin-5-yl)oxy]ethyl}-2-hydroxyacetamide;
- N*¹-{2-[(4-{[3-chloro-4-(pyridin-2-ylmethoxy)phenyl]amino}quinazolin-5-yl)oxy]ethyl}-*N*²,*N*²-dimethylglycinamide;
- N*-{2-[(4-{[3-chloro-4-(pyridin-2-ylmethoxy)phenyl]amino}quinazolin-5-yl)oxy]ethyl}-2-methoxyacetamide;
- N*-{2-[(4-{[3-chloro-4-(pyridin-2-ylmethoxy)phenyl]amino}quinazolin-5-yl)oxy]ethyl}-2-(methylsulfonyl)acetamide;
- N*-{(2*S*)-2-[(4-{[3-chloro-4-(pyridin-2-ylmethoxy)phenyl]amino}quinazolin-5-yl)oxy]propyl}-*N*-methylacetamide;

- N*-{(2*S*)-2-[(4-{[3-chloro-4-(pyridin-2-ylmethoxy)phenyl]amino}quinazolin-5-yl)oxy]propyl}-2-hydroxy-*N*-methylacetamide;
- N*¹-(2*S*)-2-[(4-{[3-chloro-4-(pyridin-2-ylmethoxy)phenyl]amino}quinazolin-5-yl)oxy]propyl)-*N*¹,*N*²,*N*²-trimethylglycinamide;
- N*-{(2*S*)-2-[(4-{[3-chloro-4-(pyridin-2-ylmethoxy)phenyl]amino}quinazolin-5-yl)oxy]propyl}-2-methoxy-*N*-methylacetamide;
- N*-{(2*S*)-2-[(4-{[3-chloro-4-(pyridin-2-ylmethoxy)phenyl]amino}quinazolin-5-yl)oxy]propyl}-*N*-methyl-2-(methylsulfonyl)acetamide;
- N*-{(2*R*)-2-[(4-{[3-chloro-4-(pyrazin-2-ylmethoxy)phenyl]amino}quinazolin-5-yl)oxy]propyl}-*N*-methylacetamide;
- N*-{(2*R*)-2-[(4-{[3-chloro-4-(1,3-thiazol-4-ylmethoxy)phenyl]amino}quinazolin-5-yl)oxy]propyl}-*N*-methylacetamide;
- N*-{(2*R*)-2-[(4-{[3-chloro-4-[(3-fluorobenzyl)oxy]phenyl]amino}quinazolin-5-yl)oxy]propyl}-*N*-methylacetamide;
- N*-{(2*R*)-2-[(4-{[3-chloro-4-[(2-fluorobenzyl)oxy]phenyl]amino}quinazolin-5-yl)oxy]propyl}-*N*-methylacetamide;
- N*-{(1*R*)-2-[(4-{[3-chloro-4-(pyridin-2-ylmethoxy)phenyl]amino}quinazolin-5-yl)oxy]-1-methylethyl}-2-hydroxy-*N*-methylacetamide;
- N*-{(1*R*)-2-[(4-{[3-chloro-4-(pyridin-2-ylmethoxy)phenyl]amino}quinazolin-5-yl)oxy]-1-methylethyl}-*N*-methylacetamide;
- N*-{(1*S*)-2-[(4-{[3-chloro-4-(pyridin-2-ylmethoxy)phenyl]amino}quinazolin-5-yl)oxy]-1-methylethyl}-2-hydroxy-*N*-methylacetamide;
- N*-{(1*S*)-2-[(4-{[3-chloro-4-(pyridin-2-ylmethoxy)phenyl]amino}quinazolin-5-yl)oxy]-1-methylethyl}-*N*-methylacetamide;
- N*-{(1*S*)-2-[(4-{[3-chloro-4-(pyridin-2-ylmethoxy)phenyl]amino}quinazolin-5-yl)oxy]-1-methylethyl}-2-methoxy-*N*-methylacetamide;
- N*-{(1*S*)-2-[(4-{[3-chloro-4-(pyridin-2-ylmethoxy)phenyl]amino}quinazolin-5-yl)oxy]-1-methylethyl}-2-hydroxyacetamide;

- N*-{(1*S*)-2-[(4-{[3-chloro-4-(pyridin-2-ylmethoxy)phenyl]amino}quinazolin-5-yl)oxy]-1-methylethyl}acetamide;
- N*¹-{(1*S*)-2-[(4-{[3-chloro-4-(pyridin-2-ylmethoxy)phenyl]amino}quinazolin-5-yl)oxy]-1-methylethyl}-*N*²,*N*²-dimethylglycinamide;
- N*¹-{(2*R*)-2-[(4-{[3-chloro-4-(pyridin-2-ylmethoxy)phenyl]amino}quinazolin-5-yl)oxy]propyl}-*N*²,*N*²-dimethylglycinamide;
- (2*S*)-*N*-{2-[(4-{[3-chloro-4-(pyridin-2-ylmethoxy)phenyl]amino}quinazolin-5-yl)oxy]ethyl}-2,4-dihydroxybutanamide;
- (2*R*)-*N*-{2-[(4-{[3-chloro-4-(pyridin-2-ylmethoxy)phenyl]amino}quinazolin-5-yl)oxy]ethyl}-2,4-dihydroxybutanamide;
- (2*R*)-*N*-{(2*R*)-2-[(4-{[3-chloro-4-(pyridin-2-ylmethoxy)phenyl]amino}quinazolin-5-yl)oxy]propyl}-2,4-dihydroxybutanamide;
- (2*S*)-*N*-{(2*R*)-2-[(4-{[3-chloro-4-(pyridin-2-ylmethoxy)phenyl]amino}quinazolin-5-yl)oxy]propyl}-2,4-dihydroxybutanamide;
- (2*R*)-*N*-{(2*S*)-2-[(4-{[3-chloro-4-(pyridin-2-ylmethoxy)phenyl]amino}quinazolin-5-yl)oxy]propyl}-2,4-dihydroxybutanamide;
- (2*S*)-*N*-{(2*S*)-2-[(4-{[3-chloro-4-(pyridin-2-ylmethoxy)phenyl]amino}quinazolin-5-yl)oxy]propyl}-2,4-dihydroxybutanamide;
- (2*S*)-*N*-{(1*R*)-2-[(4-{[3-chloro-4-(pyridin-2-ylmethoxy)phenyl]amino}quinazolin-5-yl)oxy]-1-methylethyl}-2,4-dihydroxybutanamide;
- (2*R*)-*N*-{(1*R*)-2-[(4-{[3-chloro-4-(pyridin-2-ylmethoxy)phenyl]amino}quinazolin-5-yl)oxy]-1-methylethyl}-2,4-dihydroxybutanamide;
- (2*R*)-*N*-{2-[(4-{[3-chloro-4-(1,3-thiazol-4-ylmethoxy)phenyl]amino}quinazolin-5-yl)oxy]ethyl}-2,4-dihydroxybutanamide;
- (2*S*)-*N*-{2-[(4-{[3-chloro-4-(1,3-thiazol-4-ylmethoxy)phenyl]amino}quinazolin-5-yl)oxy]ethyl}-2,4-dihydroxybutanamide;
- (2*R*)-*N*-{(1*R*)-2-[(4-{[3-chloro-4-(1,3-thiazol-4-ylmethoxy)phenyl]amino}quinazolin-5-yl)oxy]-1-methylethyl}-2,4-dihydroxybutanamide;

(2*S*)-*N*-{(1*R*)-2-[(4-{[3-chloro-4-(1,3-thiazol-4-ylmethoxy)phenyl]amino} quinazolin-5-yl)oxy]-1-methylethyl}-2,4-dihydroxybutanamide;

N-methyl-*N*-{2-[(4-{[3-methyl-4-(pyridin-2-ylmethoxy)phenyl]amino} quinazolin-5-yl)oxy]ethyl}acetamide;

N-methyl-*N*-{2-[(4-{[3-methyl-4-(1,3-thiazol-4-ylmethoxy)phenyl]amino} quinazolin-5-yl)oxy]ethyl}acetamide;

N-methyl-*N*-(2-{[4-({3-methyl-4-[(5-methylisoxazol-3-yl)methoxy]phenyl} amino} quinazolin-5-yl)oxy}ethyl)acetamide;

2-hydroxy-*N*-methyl-*N*-{2-[(4-{[3-methyl-4-(1,3-thiazol-2-ylmethoxy)phenyl]amino} quinazolin-5-yl)oxy]ethyl}acetamide;

2-hydroxy-*N*-{2-[(4-{[3-methyl-4-(pyridin-2-ylmethoxy)phenyl]amino} quinazolin-5-yl)oxy]ethyl}acetamide;

2-hydroxy-*N*-{2-[(4-{[3-methyl-4-(1,3-thiazol-4-ylmethoxy)phenyl]amino} quinazolin-5-yl)oxy]ethyl}acetamide;

N-{2-[(4-{[3-chloro-4-(pyridin-2-ylmethoxy)phenyl]amino} quinazolin-5-yl)oxy]-1,1-dimethylethyl}-2-hydroxyacetamide;

2-hydroxy-*N*-{(2*R*)-2-[(4-{[3-methyl-4-(pyridin-2-ylmethoxy)phenyl]amino} quinazolin-5-yl)oxy]propyl}acetamide;

2-hydroxy-*N*-{(2*R*)-2-[(4-{[3-methyl-4-(1,3-thiazol-4-ylmethoxy)phenyl]amino} quinazolin-5-yl)oxy]propyl}acetamide;

N-((2*R*)-2-{[4-({4-[(3-fluorobenzyl)oxy]-3-methylphenyl} amino)quinazolin-5-yl]oxy}propyl)-2-hydroxyacetamide;

2-hydroxy-*N*-{(2*R*)-2-[(4-{[3-methyl-4-(1,3-thiazol-2-ylmethoxy)phenyl]amino} quinazolin-5-yl)oxy]propyl}acetamide;

N-{(2*R*)-2-[(4-{[3-methyl-4-(pyridin-2-ylmethoxy)phenyl]amino} quinazolin-5-yl)oxy]propyl}acetamide;

N-{(2*R*)-2-[(4-{[3-methyl-4-(1,3-thiazol-4-ylmethoxy)phenyl]amino} quinazolin-5-yl)oxy]propyl}acetamide;

N-((2*R*)-2-{{4-({4-(3-fluorobenzyl)oxy}-3-methylphenyl)amino}quinazolin-5-yl}oxy)propyl)acetamide;

N-{(2*R*)-2-[(4-{{3-methyl-4-(1,3-thiazol-2-ylmethoxy)phenyl}amino}quinazolin-5-yl)oxy]propyl}acetamide;

2-hydroxy-*N*-methyl-*N*-{(2*R*)-2-[(4-{{3-methyl-4-(pyridin-2-ylmethoxy)phenyl}amino}quinazolin-5-yl)oxy]propyl}acetamide;

2-hydroxy-*N*-methyl-*N*-{(2*R*)-2-[(4-{{3-methyl-4-(1,3-thiazol-4-ylmethoxy)phenyl}amino}quinazolin-5-yl)oxy]propyl}acetamide;

2-hydroxy-*N*-methyl-*N*-{(2*R*)-2-[(4-{{3-methyl-4-((5-methylisoxazol-3-yl)methoxy)phenyl}amino}quinazolin-5-yl)oxy]propyl}acetamide;

N-methyl-*N*-{(1*R*)-1-methyl-2-[(4-{{3-methyl-4-(pyridin-2-ylmethoxy)phenyl}amino}quinazolin-5-yl)oxy]ethyl}acetamide;

N-methyl-*N*-{(1*R*)-1-methyl-2-[(4-{{3-methyl-4-(1,3-thiazol-4-ylmethoxy)phenyl}amino}quinazolin-5-yl)oxy]ethyl}acetamide;

N-{(1*R*)-2-[(4-{{3-chloro-4-(1,3-thiazol-4-ylmethoxy)phenyl}amino}quinazolin-5-yl)oxy]-1-methylethyl}-2-hydroxy-*N*-methylacetamide;

2-hydroxy-*N*-methyl-*N*-{(1*R*)-1-methyl-2-[(4-{{3-methyl-4-(pyridin-2-ylmethoxy)phenyl}amino}quinazolin-5-yl)oxy]ethyl}acetamide;

2-hydroxy-*N*-methyl-*N*-{(1*R*)-1-methyl-2-[(4-{{3-methyl-4-(1,3-thiazol-4-ylmethoxy)phenyl}amino}quinazolin-5-yl)oxy]ethyl}acetamide;

N-{(2*R*)-2-[(4-{{3-chloro-4-(pyridin-2-ylmethoxy)phenyl}amino}quinazolin-5-yl)oxy]propyl}-1-hydroxy-*N*-methylcyclopropanecarboxamide;

(2*S*)-*N*-{(2*R*)-2-[(4-{{3-chloro-4-(pyridin-2-ylmethoxy)phenyl}amino}quinazolin-5-yl)oxy]propyl}-2-hydroxy-*N*-methylpropanamide;

N-{(2*R*)-2-[(4-{{3-chloro-4-(pyridin-2-ylmethoxy)phenyl}amino}quinazolin-5-yl)oxy]propyl}-2-hydroxy-*N*,2-dimethylpropanamide;

(2*R*)-*N*-{(2*R*)-2-[(4-{{3-chloro-4-(pyridin-2-ylmethoxy)phenyl}amino}quinazolin-5-yl)oxy]propyl}-2-hydroxy-*N*-methylpropanamide;

(2*R*)-*N*-{(2*R*)-2-[(4-{[3-chloro-4-(pyridin-2-ylmethoxy)phenyl]amino}quinazolin-5-yl)oxy]propyl}-2-methoxy-*N*-methylpropanamide;

2-hydroxy-*N*-methyl-*N*-((2*R*)-2-{[4-({3-methyl-4-[(6-methylpyridin-3-yl)oxy]phenyl} amino)quinazolin-5-yl]oxy}propyl)acetamide;

N-methyl-*N*-((2*R*)-2-{[4-({3-methyl-4-[(6-methylpyridin-3-yl)oxy]phenyl} amino)quinazolin-5-yl]oxy}propyl)acetamide;

*N*¹,*N*²,*N*³-trimethyl-*N*¹-((2*R*)-2-{[4-({3-methyl-4-[(6-methylpyridin-3-yl)oxy]phenyl} amino)quinazolin-5-yl]oxy}propyl)glycinamide;

N-methyl-*N*-((2*R*)-2-{[4-({3-methyl-4-[(6-methylpyridin-3-yl)oxy]phenyl} amino)quinazolin-5-yl]oxy}propyl)-2-pyrrolidin-1-ylacetamide;

N-methyl-*N*-((2*R*)-2-{[4-({3-methyl-4-[(6-methylpyridin-3-yl)oxy]phenyl} amino)quinazolin-5-yl]oxy}propyl)-2-morpholin-4-ylacetamide;

N-methyl-*N*-((2*R*)-2-{[4-({3-methyl-4-[(6-methylpyridin-3-yl)oxy]phenyl} amino)quinazolin-5-yl]oxy}propyl)-2-(4-methylpiperazin-1-yl)acetamide;

2-hydroxy-*N*-methyl-*N*-((2*S*)-2-{[4-({3-methyl-4-[(6-methylpyridin-3-yl)oxy]phenyl} amino)quinazolin-5-yl]oxy}propyl)acetamide;

N-methyl-*N*-((2*S*)-2-{[4-({3-methyl-4-[(6-methylpyridin-3-yl)oxy]phenyl} amino)quinazolin-5-yl]oxy}propyl)acetamide;

N-methyl-*N*-((2*S*)-2-{[4-({3-methyl-4-[(6-methylpyridin-3-yl)oxy]phenyl} amino)quinazolin-5-yl]oxy}propyl)-2-pyrrolidin-1-ylacetamide;

(2*S*)-2,4-dihydroxy-*N*-((2*R*)-2-{[4-({3-methyl-4-[(6-methylpyridin-3-yl)oxy]phenyl} amino)quinazolin-5-yl]oxy}propyl)butanamide;

(2*S*)-4-bromo-2-hydroxy-*N*-((2*R*)-2-{[4-({3-methyl-4-[(6-methylpyridin-3-yl)oxy]phenyl} amino)quinazolin-5-yl]oxy}propyl)butanamide;

N-(2-chloroethyl)-*N*-((2*R*)-2-{[4-({3-methyl-4-[(6-methylpyridin-3-yl)oxy]phenyl} amino)quinazolin-5-yl]oxy}propyl)urea;

2-hydroxy-*N*-methyl-*N*-((1*R*)-1-methyl-2-{{4-({3-methyl-4-[(6-methylpyridin-3-yl)oxy]phenyl} amino)quinazolin-5-yl}oxy} ethyl)acetamide;

N-methyl-*N*-((1*R*)-1-methyl-2-{{4-({3-methyl-4-[(6-methylpyridin-3-yl)oxy]phenyl} amino)quinazolin-5-yl}oxy} ethyl)acetamide;

2-hydroxy-*N*-methyl-*N*-((1*S*)-1-methyl-2-{{4-({3-methyl-4-[(6-methylpyridin-3-yl)oxy]phenyl} amino)quinazolin-5-yl}oxy} ethyl)acetamide;

N-methyl-*N*-((1*S*)-1-methyl-2-{{4-({3-methyl-4-[(6-methylpyridin-3-yl)oxy]phenyl} amino)quinazolin-5-yl}oxy} ethyl)acetamide;

methyl- {2-[(4- { [3-chloro-4-(pyridin-2-ylmethoxy)phenyl]amino} quinazolin-5-yl)oxy]ethyl} methylcarbamate;

N- {2-[(4- { [3-chloro-4-(pyridin-2-ylmethoxy)phenyl]amino} quinazolin-5-yl)oxy]ethyl} -*N,N*-dimethylurea;

N-(2-chloroethyl)-*N*- {2-[(4- { [3-chloro-4-(pyridin-2-ylmethoxy)phenyl]amino} quinazolin-5-yl)oxy]ethyl} -*N*-methylurea;

N- { (2*R*)-2-[(4- { [3-chloro-4-(pyridin-2-ylmethoxy)phenyl]amino} quinazolin-5-yl)oxy]propyl} -*N*-methylurea;

[[(1*R*)-2- {4- [3-chloro-4-(pyridin-2-ylmethoxy)phenyl]amino} quinazolin-5-yl)oxy]propylcarbamoyl)methyl]methylcarbamic acid tert-butyl ester;

*N*¹- { (2*R*)-2-[(4- { [3-chloro-4-(pyridin-2-ylmethoxy)phenyl]amino} quinazolin-5-yl)oxy]propyl} -*N*²-methylglycinamide;

2-hydroxy-*N*-methyl-*N*-(2-{{4-({3-methyl-4-[(6-methylpyridin-3-yl)oxy]phenyl} amino)quinazolin-5-yl}oxy} ethyl)acetamide;

N-methyl-*N*-(2-{{4-({3-methyl-4-[(6-methylpyridin-3-yl)oxy]phenyl} amino)quinazolin-5-yl}oxy} ethyl)acetamide; and

N- {2-[(4- { [3-chloro-4-(1-methyl-1-pyridin-2-ylethoxy)phenyl]amino} quinazolin-5-yl)oxy]ethyl} -*N*-methylacetamide;


or a pharmaceutically acceptable salt thereof.

Claim 19 (previously presented): A pharmaceutical composition which comprises a quinazoline derivative of the formula I, or a pharmaceutically acceptable salt thereof, as defined in claim 1 or claim 18 in association with a pharmaceutically-acceptable diluent or carrier.

Claims 20-23 (cancelled).

Claim 24 (currently amended): A process for the preparation of preparing a quinazoline derivative of the formula I, or a pharmaceutically acceptable salt thereof, as defined in claim 1 which comprises:

- a) ~~the~~ coupling, optionally in the presence of a suitable base, of a quinazoline of the formula II:



II


wherein R^1 , R^2 , R^3 , R^4 , R^{4a} , R^5 , R^{5a} , R^6 , X^1 , Q^1 , m , and n have any of the meanings defined in claim 1 except that any functional group is optionally protected, with a carboxylic acid of the formula III, or a reactive derivative thereof:



III

wherein A has any of the meanings defined in claim 1 except that any functional group is optionally protected if necessary; or

- (b) for the preparation of ~~these the~~ compounds of the formula I wherein X^1 is $OC(R^7)_2$, $SC(R^7)_2$ or $N(R^7)C(R^7)_2$, ~~the reaction reacting, conveniently optionally~~ in the presence of a suitable base, of a quinazoline of the formula IV:



IV

wherein X^{1a} is O, S or $N(R^7)_2$; and R^1 , R^2 , R^3 , R^4 , R^{4a} , R^5 , R^{5a} , R^6 , R^7 , A, m, and n have any of the meanings defined in claim 1 except that any functional group is optionally protected, with a compound of the formula V or a salt thereof:



V

wherein L^1 is a suitable displaceable group and Q^1 and R^7 have any of the meanings defined in claim 1 except that any functional group is optionally protected ~~if necessary; or~~


- (c) for the preparation of ~~these the~~ compounds of the formula I wherein A is R^{14} and R^{14} is NHR^{17} or Q^3-X^5 - (wherein R^{17} and Q^3 are as defined in claim 1 and X^5 is NH), the coupling of a quinazoline of the formula II as defined above in (a) with an isocyanate of the formula IIIa:



IIIa


wherein A is R^{14} as previously defined in this section except that any functional group is optionally protected; or

- (d) ~~the reaction of reacting~~ a quinazoline of the formula II wherein R^6 is hydrogen:


**II**

wherein R^1 , R^2 , R^3 , R^4 , R^{4a} , R^5 , R^{5a} , X^1 , Q^1 , m , and n have any of the meanings defined in claim 1 except that any functional group is optionally protected, with α -hydroxy- γ -butyrolactone wherein any functional group is optionally protected; or

(c) the coupling of a quinazoline of the formula VI:


**VI**

wherein R^1 , R^4 , R^{4a} , R^5 , R^{5a} , R^6 , A and m have any of the meanings defined in claim 1 except that any functional group is optionally protected, with a compound of the formula IIb:

**IIb**

wherein R^2 , R^3 , X^1 , Q^1 and n have any of the meanings defined in claim 1 except that any functional group is optionally protected; or


- (f) for the preparation of ~~these the~~ compounds of the formula I wherein X^1 is O and Q^1 is 2-pyridyl, 4-pyridyl, 2-pyrimidyl, 4-pyrimidyl, 2-pyrazinyl or 3-pyridazinyl, ~~the reaction, reacting, conveniently optionally~~ in the presence of a suitable base and a suitable catalyst, of a quinazoline of the formula VII:



VII

wherein R^1 , R^2 , R^3 , R^4 , R^{4a} , R^5 , R^{5a} , R^6 , A, m and n have any of the meanings defined in claim 1 except that any functional group is optionally protected, with 2-bromopyridine, 4-bromopyridine, 2-chloropyrimidine, 4-chloropyrimidine, 2-chloropyrazine or 3-chloropyridazine; or

- (g) for the preparation of ~~these the~~ compounds of the formula I wherein A is $Z-(CR^{12}R^{13})_p$, wherein Z is $NR^{16}R^{17}$, the reaction, ~~conveniently optionally~~ in the presence of a suitable base, of a quinazoline of the formula VIII:



VIII

wherein L^1 is a suitable displaceable group and $R^1, R^2, R^3, R^4, R^{4a}, R^5, R^{5a}, R^6, R^{12}, R^{13}, X^1, Q^1, m, n$ and p have any of the meanings defined in claim 1 except that any functional group is optionally protected, with a compound of the formula **IXa**, or a reactive derivative thereof:



IXa

wherein R^{16} and R^{17} have any of the meanings defined in claim 1 except that any functional group is optionally protected;

and thereafter, optionally:

- (i) converting a quinazoline derivative of the formula I into another quinazoline derivative of the formula I;
- (ii) removing any protecting group that is present;
- (iii) forming a pharmaceutically acceptable salt.

Claim 25 (currently amended): A method for treating a breast tumour sensitive to inhibition of an ~~erbB2 receptor tyrosine kinase~~ in a warm-blooded animal in need of such treatment, which comprises administering to ~~said the~~ animal an effective amount of a quinazoline derivative of the formula I, or a pharmaceutically acceptable salt thereof, according to claim 1.

Claims 26-30 (cancelled).